

Figure 1B

Figure 2a




Figure 2b




Figure 3





Figure 4






Figure 5




Figure 6




Figure 7

Figure 7 shows the relationship between NTN concentration and ELISA (Absorbance) and Immuno-PCR (Ct) values. The x-axis represents NTN concentration in pg/ml on a logarithmic scale from 0.01 to 10<sup>5</sup>. The left y-axis represents ELISA (Absorbance) from 0 to 2. The right y-axis represents Immuno-PCR (Ct) from 14 to 30. Two data series are plotted: Absorbance (squares) and Ct (circles). Both series show a decreasing trend as NTN concentration increases. The Absorbance series starts at approximately 1.7 at 0.01 pg/ml and decreases to approximately 0.3 at 10<sup>5</sup> pg/ml. The Ct series starts at approximately 28.5 at 0.01 pg/ml and decreases to approximately 17.5 at 10<sup>5</sup> pg/ml. Error bars are present for the Ct data points at 1000, 10000, and 100000 pg/ml.






Figure 8

bioRxiv preprint doi: <https://doi.org/10.1101/111111>; this version posted January 1, 2017. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.